

Microsphere assays used to detect plant pathogens

01 July 2013 | News | By BioSpectrum Bureau



Singapore: Thai researchers have developed a novel multiplex detection method based on microsphere immunoassays to simultaneously detect four important plant pathogens, including, a fruit blotch bacterium *Acidovorax avenae* subsp. *citrulli* (Aac); chilli vein-banding mottle virus (CVbMV, potyvirus), watermelon silver mottle virus (WSMoV, tospovirus serogroup IV) and melon yellow spot virus (MYSV, tospovirus).

The paper, describing the development of a multiplex microsphere immunoassay to detect the widespread plant pathogens, was published in Plos One in April 2013. The work is a collaborative effort between scientists from BIOTEC and Queen's University Belfast, UK.

An antibody for each plant pathogen was linked on a fluorescence-coded magnetic microsphere set which was used to capture corresponding pathogen. The presence of pathogens was detected by R-phycoerythrin (RPE)-labeled antibodies specific to the pathogens. The assay was able to detect all four plant pathogens precisely and accurately with substantially higher sensitivity than enzyme-linked immunosorbent assay (ELISA).

The assay time of the microsphere immunoassay is an one-hour, as opposed to four-hours required by that of ELISA. This system was also shown to be capable of detecting the pathogens in naturally infected plant samples.

The capacity of this microsphere immunoassay technique could be further expanded to higher throughput such as detecting up to 50 targets simultaneously. The system could also become fully automatic if dealing with a larger volume of routine testing.